

前立腺がん検診受診者における男性型脱毛症治療薬（AGA）の服用について

資料11

令和6年度 前立腺がん検診受診者 (n=7,396)

結果判定（要精検者・率）

区分	受診者数	結果判定			
		要精検		精検不要	
50歳	1,303	47	3.6%	1,256	96.4%
55歳	1,502	75	5.0%	1,427	95.0%
60歳	1,504	146	9.7%	1,358	90.3%
65歳	1,635	143	8.7%	1,492	91.3%
70歳	1,452	147	10.1%	1,305	89.9%
計	7,396	558	7.5%	6,838	92.5%

AGA内服の有無

区分	受診者数	AGA内服					
		あり		なし		空白	
50歳	1,303	148	11.4%	1,149	88.2%	6	0.5%
55歳	1,502	138	9.2%	1,357	90.3%	7	0.5%
60歳	1,504	113	7.5%	1,380	91.8%	11	0.7%
65歳	1,635	106	6.5%	1,524	93.2%	5	0.3%
70歳	1,452	69	4.8%	1,377	94.8%	6	0.4%
計	7,396	574	7.8%	6,787	91.8%	35	0.5%

AGA内服あり (n=574)

区分	受診者数	AGA 内服あり	結果判定						
			要精検		精検不要		(再掲) PSA2倍値で要精検		
50歳	1,303	148	11.4%	2	1.4%	146	98.6%	14	9.5%
55歳	1,502	138	9.2%	4	2.9%	134	97.1%	8	5.8%
60歳	1,504	113	7.5%	7	6.2%	106	93.8%	17	15.0%
65歳	1,635	106	6.5%	6	5.7%	100	94.3%	12	11.3%
70歳	1,452	69	4.8%	6	8.7%	63	91.3%	7	10.1%
計	7,396	574	7.8%	25	4.4%	549	95.6%	58	10.1%

要精検者25名は、全員、検診のPSA測定値が判定基準の要精検の基準値内。

AGA内服なし (n=6,787)

区分	受診者数	AGA 内服なし	結果判定				
			要精検		精検不要		
50歳	1,303	1,149	88.2%	45	3.9%	1,104	96.1%
55歳	1,502	1,357	90.3%	71	5.2%	1,286	94.8%
60歳	1,504	1,380	91.8%	137	9.9%	1,243	90.1%
65歳	1,635	1,524	93.2%	137	9.0%	1,387	91.0%
70歳	1,452	1,377	94.8%	140	10.2%	1,237	89.8%
計	7,396	6,787	91.8%	530	7.8%	6,257	92.2%